



Zinc Sulfate

Injection, USP

FREQUENTLY ASKED QUESTIONS

Pack NDC#	Strength	Supplied as	Shelf pack
0517-6101-25	1 mg/mL	10 mL Pharmacy bulk package vial	25
0517-6103-25	3 mg/mL	10 mL Pharmacy bulk package vial	25
0517-8005-25	5 mg/mL	5 mL Pharmacy bulk package vial	25

- 1. I use an automated compounding device for parenteral nutrition (PN) preparations. Are there differences in the specific gravity, osmolarity, and any other intrinsic values that I need to know to program into my compounding device?**

For Zinc Sulfate Injection, USP, the following values apply:

	1 mg/mL	3 mg/mL	5 mg/mL
Specific gravity	1.003 g/mL	1.008 g/mL	1.014 g/mL
Osmolarity	33 mOsmol/L	96.5 mOsmol/L	157.2 mOsmol/L
pH range ¹	2 to 4	2 to 4	2 to 4

- 2. How should Zinc Sulfate Injection, USP be administered?**

Zinc Sulfate Injection, USP is for *admixture use* only. It is *not for direct intravenous infusion*. Prior to administration, Zinc Sulfate Injection, USP must be transferred to a separate PN container, diluted, and used as an admixture in PN solutions.¹

- 3. What is the difference between a single-dose vial (SDV) and a pharmacy bulk package (PBP) vial?**

- An SDV is intended for single use
- A PBP vial is a sterile preparation that contains many single doses

- 4. How often can the PBP vial be penetrated?**

Penetrate vial closure only one time with a suitable sterile transfer device or a dispensing set that allows measured dispensing of the contents.

- 5. How stable is PBP-prepared Zinc Sulfate Injection, USP?**

- Use Zinc Sulfate Injection, USP for admixing promptly once the sterile transfer set has been inserted into the PBP vial, or not more than 4 hours at room temperature (20°C to 25°C or 68°F or 77°F) after the vial closure has been penetrated. Discard any remaining admixture¹

- Use PN solution containing Zinc Sulfate Injection, USP promptly after mixing. Any storage of the admixture should be under refrigeration from 2°C to 8°C (36°F to 46°F) and limited to a period of time no longer than 9 days. After removal from refrigeration, use promptly and complete the infusion within 24 hours. Discard any remaining admixture¹
- Protect the admixed PN solution from light¹

6. Are the Zinc Sulfate Injection, USP formulations by American Regent, Inc.® preservative-free?

The Zinc Sulfate Injection, USP formulations are preservative-free.¹

7. Is Zinc Sulfate latex-free?

The vial closures are not made with natural rubber latex.

Zinc Sulfate

Injection, USP

For intravenous use

INDICATIONS AND USAGE

Zinc Sulfate is a trace element indicated in adult and pediatric patients as a source of zinc for parenteral nutrition when oral or enteral nutrition is not possible, insufficient, or contraindicated.

IMPORTANT ADMINISTRATION INFORMATION

Zinc Sulfate Injection is supplied as a pharmacy bulk package for *admixing use* only. It is *not for direct intravenous infusion*.

IMPORTANT SAFETY INFORMATION

CONTRAINDICATIONS

Zinc Sulfate Injection is contraindicated in patients with known hypersensitivity to zinc.

WARNINGS AND PRECAUTIONS

Pulmonary Embolism due to Pulmonary Vascular Precipitates:

If signs of pulmonary distress occur, stop the infusion and initiate a medical evaluation. The infusion set and catheter should be checked periodically for precipitates.

Vein Damage and Thrombosis: Zinc Sulfate Injection has a low pH and must be prepared and used as an admixture in PN solutions. Solutions with osmolarity of 900 mOsm/L or more must be infused through a central venous catheter.

Aluminum Toxicity: Zinc Sulfate Injection contains aluminum that may be toxic. Aluminum may reach toxic levels with prolonged parenteral administration if kidney function is impaired. Preterm infants are particularly at risk for aluminum toxicity because their kidneys are immature, and they require large amounts of calcium and phosphate solutions, which also contain aluminum.

Monitoring and Laboratory Tests: Monitor Zinc concentrations, fluid and electrolyte status, serum osmolarity, blood glucose, liver and kidney function, blood count and coagulation parameters throughout treatment.

Copper Deficiency: Several post-marketing cases have reported that high doses of supplemental zinc (approximately 10 times the recommended dosage of 3 mg/day Zinc Sulfate Injection in adults) taken over extended periods of time (ie, months to years) may result in decreased enteral copper absorption and copper deficiency.

Hypersensitivity Reactions: If hypersensitivity reactions occur, discontinue Zinc Sulfate Injection and initiate appropriate medical treatment.

ADVERSE REACTIONS

No zinc-related adverse reactions have been reported in clinical studies or postmarketing reports in patients receiving intravenously administered PN solutions containing zinc sulfate within the recommended dosage range.

USE IN SPECIFIC POPULATIONS

Pregnancy: *Risk Summary:* Administration of the recommended dose of Zinc Sulfate Injection in PN is not expected to cause major birth defects, miscarriage, or adverse maternal or fetal outcomes.

Lactation: *Risk Summary:* Zinc is present in human milk. There is no information on the effects of zinc sulfate on milk production. The developmental and health benefits of breastfeeding should be considered along with the mother's clinical need for Zinc Sulfate Injection and any potential adverse effects on the breastfed infant from Zinc Sulfate Injection or from the underlying maternal condition.

Pediatric Use: Safety and dosing recommendations in pediatric patients are based on published literature describing controlled studies of zinc-containing products in pediatric patients.

Geriatric Use: Dose selection should be individualized based on the patient's clinical condition, nutritional requirements, and additional nutritional intake provided orally or enterally to the patient.

OVERDOSAGE: There are reported cases of overdosage with intravenous zinc in parenteral nutrition.

For additional safety information, please see [Full Prescribing Information](#).

You are encouraged to report adverse drug events to American Regent, Inc. at 1-800-734-9236 or to the FDA by visiting www.fda.gov/medwatch or calling 1-800-FDA-1088.

You are encouraged to report adverse drug events (ADEs) to American Regent®:

T 1.800.734.9236; **E** pv@americanregent.com; **F** 1.610.650.0170;

ADEs may also be reported to the FDA:

1.800.FDA.1088

or www.fda.gov/medwatch

Medical Information:

T 1.888.354.4855 (9:00 am–5:00 pm Eastern Time, Monday–Friday)
www.americanregent.com/medical-affairs

For medical information outside of normal business hours
that cannot wait until the next business day, please call 1.877.845.6371

REFERENCE:

1. Zinc Sulfate Injection, USP. Package Insert. American Regent, Inc.

5 Ramsey Road \ Shirley, NY 11967 \ 1.800.645.1706
F 631.924.1731 \ www.americanregent.com

© 2023 American Regent, Inc. All rights reserved. PP-ZN-US-0011 v3.0

